

JOURNAL OF ALGEBRA 74, 211–222 (1982)

Duality and Character Values of Finite Groups of Lie Type

DEAN ALVIS

*Department of Mathematics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139*

Communicated by W. Feit

Received March 12, 1981

INTRODUCTION

The purpose of this paper is to show how the values of the dual of a character of a finite group of Lie type can be computed on semisimple classes, given sufficient information about the original character.

In the first approach, the values of a character χ on elements su with (fixed) semisimple part s are related to the value of the dual χ^* at s . To be specific, $\sum \chi(su) = \text{St}(s) \chi^*(s)$, where St is the Steinberg character. The special case in which χ is a component of the permutation character 1_B^G and $s = 1$ was established by Kawanaka in [15].

A second method can be used for the group of rational points of a connected, reductive algebraic group defined over a finite field with q elements. It is shown that $\chi^*(s) = \text{St}(s) \chi^*(s)$, where, roughly speaking, $\chi^*(s)$ is $\chi(s)$ with q replaced by $1/q$ (see Section 3). When $s = 1$ and χ is a component of 1_B^G , this formula coincides with a well-known formula involving generic degrees (see Green [12] and Curtis [9]).

In Section 4, both approaches are applied to prove, for large q , a conjecture of I. Macdonald (see [21, 6.11]). Special cases of this conjecture have been proved in Kawanaka [15] and Springer [22].

Most of the results in this paper were announced in [2] and proved in the author's dissertation [3].

1. PRELIMINARY RESULTS

If H is a finite group, $\text{irr } H$ will denote the set of irreducible characters of H , $\text{char } H$ the virtual characters of H , and $\text{cf } H$ the class functions on H , all over the field \mathbb{C} of complex numbers. Let $(\ , \)_H$, or simply $(\ , \)$, denote the usual Hermitian form on $\text{cf } H$; $(f, g) = |H|^{-1} \sum_{h \in H} f(h) \overline{g(h)}$.

Let G be a finite group with split (B, N) -pair of characteristic p (see [6, 8, 20]), $G = \langle B, N \rangle$. Let (W, R) be the Coxeter system for G ; R is a set of distinguished generators for the Coxeter group $W = N/T$, $T = B \cap B$. For $J \subseteq R$ let W_J be the subgroup of W generated by J . Then $P_J = BW_JB$ is a standard parabolic subgroup of G , with Levi decomposition $P_J = L_J V_J$, $V_J = O_p(P_J)$. The Levi factor L_J has a split (B, N) -pair of characteristic p , with Coxeter system (W_J, J) . If $K \subseteq J$, then $P_{J,K} = L_J \cap P_K$ is a standard parabolic subgroup of L_J , with Levi decomposition $P_{J,K} = L_K V_{J,K}$, $V_{J,K} = L_J \cap V_K$. Also, V_K is the semidirect product $V_K = V_J V_{J,K}$.

For $K \subseteq J \subseteq R$ and $\varphi \in \text{irr } L_K$, let $\tilde{\varphi}$ be the extension of φ to $P_{J,K}$ via the projection $P_{J,K} \rightarrow L_K \cong P_{J,K}/V_{J,K}$. Thus $\tilde{\varphi}(lv) = \varphi(l)$ for $l \in L_K$, $v \in V_{J,K}$. We define $I_K^J \varphi = \text{Ind}_{P_{J,K}}^{L_J} \tilde{\varphi}$. Note that I_K^J extends to linear maps $I_K^J: \text{char } L_K \rightarrow \text{char } L_J$ and $I_K^J: \text{cf } L_K \rightarrow \text{cf } L_J$. We will write I_K instead of I_K^R when $J = R$.

Assume now that $K \subseteq J \subseteq R$ and $\chi \in \text{irr } L_J$. Then define $T_K^J \chi = \sum_{\varphi \in \text{irr } L_K} (\chi, I_K^J \varphi)_{L_J} \varphi$; $T_K^J \chi$ is called the truncation of χ to L_K (Curtis [9]). We will write T_K instead of T_K^R when $J = R$. If χ is the character of the $\mathbb{C}L_J$ -module M , then $T_K^J \chi$ is the character of L_K afforded by the subspace of $V_{J,K}$ -invariants of M . Note that T_K^J extends to linear maps $T_K^J: \text{char } L_J \rightarrow \text{char } L_K$ and $T_K^J: \text{cf } L_J \rightarrow \text{cf } L_K$. If $f \in \text{cf } L_J$, then the value of $T_K^J f$ at $l \in L_K$ is given by

$$T_K^J f(l) = |V_{J,K}|^{-1} \sum_{v \in V_{J,K}} f(lv). \quad (1.1)$$

The next two lemmas follow easily from the remarks above and will be stated without proof.

(1.2) LEMMA. *If $K \subseteq J \subseteq R$, $f \in \text{cf } L_J$ and $g \in \text{cf } L_K$, then $(f, I_K^J g)_{L_J} = (T_K^J f, g)_{L_K}$.*

(1.3) LEMMA. *If $K \subseteq J \subseteq R$, then $I_K = I_J I_K^J$ and $T_K = T_K^J T_K$.*

Let f be a class function on G . Following Curtis [9], we define the dual f^* of f to be the class function $f^* = \sum_{J \subseteq R} (-1)^{|J|} I_J T_J f$.

(1.4) THEOREM (Curtis [9, Theorem 1.3]). *If $J \subseteq R$ and $f \in \text{cf } G$, then $(T_J f)^* = T_J(f^*)$.*

The proof of the following theorem, which is outlined in [1], will be included for the sake of completeness. This theorem has also been proved independently by N. Kawanaka (private communication).

(1.5) THEOREM. *The duality map $f \rightarrow f^*$ is an isometry on $\text{cf } G$ of order two. Thus $\chi^{**} = \chi$ and $\pm \chi^* \in \text{irr } G$ if $\chi \in \text{irr } G$.*

Proof. Let $f, g \in \text{cf } G$. Using the definition for f^* , we see that (f^*, g) is equal to

$$\sum_{J \subseteq R} (-1)^{|J|} \sum_{\varphi \in \text{irr } L_J} (f, I_J \varphi) (I_J \varphi, g). \quad (1.6)$$

The symmetry of (1.6) shows that $(f^*, g) = (f, g^*)$. It remains to prove $f^{**} = f$.

We have $f^{**} = \sum_{J \subseteq R} (-1)^{|J|} I_J T_J(f^*) = \sum_{J \subseteq R} (-1)^{|J|} I_J ((T_J f)^*)$ by Curtis' result, Theorem 1.4. Expanding $(T_J f)^*$, we obtain

$$f^{**} = \sum_{J \subseteq R} (-1)^{|J|} \sum_{K \subseteq J} (-1)^{|K|} I_J I_K^J T_K^J T_J f. \quad (1.7)$$

Finally, applying Lemma 1.3 to (1.7) gives

$$f^{**} = \sum_{K \subseteq R} (-1)^{|K|} \left(\sum_{J \supseteq K} (-1)^{|J|} \right) I_K T_K f. \quad (1.8)$$

For a given $K \subseteq R$, the contribution to the sum in (1.8) is zero unless $K = R$. Therefore $f^{**} = I_R T_R f = f$, which finishes the proof of the theorem.

Let $J \subseteq R$ and assume $\varphi \in \text{irr } L_J$ is cuspidal. That is, $T_K^J \varphi = 0$ when $K \subset J$, $K \neq J$. An immediate consequence of Theorems 1.4 and 1.5 and Lemma 1.2 is that $(I_J \varphi)^* = I_J(\varphi^*) = (-1)^{|J|} I_J \varphi$. If $\chi \in \text{irr } G$ is a component of $I_J \varphi$ of multiplicity m , so that $m = (\chi, I_J \varphi) > 0$, we deduce from Theorem 1.5 that $(-1)^{|J|} \chi^* \in \text{irr } G$, and that $(-1)^{|J|} \chi^*$ is also a component of $I_J \varphi$ of multiplicity m .

2. DUALITY AND JORDAN DECOMPOSITION

Suppose H is a finite group, $x \in H$, and that p is a prime. There is a unique p -element u of H and a unique p' -element s of H such that $x = su = us$; u (s) is called the p -part (p' -part, respectively) of x . The proof of the following lemma is elementary and will be omitted (see Lusztig [17, Sect. 2]).

(2.1) LEMMA. *Let P be a finite group which is a semidirect product $P = LV$, where V is a normal p -subgroup of P . Assume $l \in L$ and $v \in V$. Then the p' -parts of l and lv are conjugate by an element of V .*

Now let $G = \langle B, N \rangle$ be a finite group with split (B, N) -pair of characteristic p . If $x \in G$, the p -part (p' -part) of x will be called the unipotent (semisimple) part of x , and will be denoted by x_{un} (x_{ss} , respectively). The

element x will be called unipotent (semisimple) if $x = x_{un}$ ($x = x_{ss}$, respectively).

It will be convenient to use a basis for $\text{cf } G$ other than $\text{irr } G$. For $x, y \in G$, define $\rho_x(y) = |Z(x)|$ if x and y are conjugate, and set $\rho_x(y) = 0$ otherwise. Then $\{\rho_x \mid x \in G\}$ is a basis for $\text{cf } G$, and $(f, \rho_x) = f(x)$ for $f \in \text{cf } G$, $x \in G$.

(2.2) LEMMA. *If $x, y \in G$ and $\rho_x^*(y) \neq 0$, then x_{ss} and y_{ss} are conjugate.*

Proof. If $\rho_x^*(y) \neq 0$, then $I_J T_J \rho_x(y) \neq 0$ for some $J \subseteq R$. Thus there are $l \in L_J$ and $v \in V_J$ such that y is conjugate to lv and $(T_J \rho_x)^{\sim}(lv) = T_J \rho_x(l) \neq 0$. From (1.1) we see that $\rho_x(lu) \neq 0$ for some $u \in V_J$, so that x is conjugate to lu . The proof is completed by applying Lemma 2.1 to $P_J = L_J V_J$.

Let $s \in G$ be semisimple. Denote by $V(s)$ the set of all unipotent elements in $Z(s)$, the centralizer of s .

(2.3) THEOREM. *Let $s \in G$ be semisimple, and assume $f, g \in \text{cf } G$. Then*

$$\sum_{u \in V(s)} f(su) g(su) = \sum_{u \in V(s)} f^*(su) g^*(su).$$

Proof. It is enough to argue in the case $f = \rho_x$, $g = \rho_y^*$, $x, y \in G$, by Theorem 1.5. It must be shown that

$$\sum_{u \in V(s)} \rho_x(su) \rho_y^*(su) = \sum_{u \in V(s)} \rho_x^*(su) \rho_y(su),$$

or, equivalently, that

$$\rho_y^*(x) \sum_{u \in V(s)} \rho_x(su) = \rho_x^*(y) \sum_{u \in V(s)} \rho_y(su). \quad (2.4)$$

By Lemma 2.2, both sides of (2.4) vanish unless x_{ss} and y_{ss} are conjugate. We may reduce to the case $x = su_0$, $y = sv_0$, $u_0, v_0 \in V(s)$. An easy calculation shows both sums in (2.4) are equal to $|Z(s)|$. Finally, $\rho_y^*(x) = (\rho_y^*, \rho_x) = (\rho_y, \rho_x^*) = (\rho_x^*, \rho_y) = \rho_x^*(y)$ by Theorem 1.5. This completes the proof of the theorem.

Let St be the Steinberg character of G . Recall that St is an irreducible character of G of degree $|G|_p$, and that $\text{St} = \sum_{J \subseteq R} (-1)^{|J|} \text{Ind}_{P_J}^G 1_{P_J}$ (Curtis [7, Theorem 2]). Thus $\text{St} = 1_G^*$, and St vanishes off the semisimple classes of G . The following three corollaries are therefore special cases of Theorem 2.3.

(2.5) COROLLARY. *Let $s \in G$ be semisimple, and let χ be a character of G . Then $\sum_{u \in V(s)} \chi(su) = \text{St}(s) \chi^*(s)$.*

(2.6) COROLLARY. *If χ is a character of G , then $\sum_{u \in V(1)} \chi(u) = |G|_p \chi^*(1)$.*

(2.7) COROLLARY (Steinberg [23]). *If $s \in G$ is semisimple, then $|V(s)| = \text{St}(s)^2$.*

The next result can be thought of as an orthogonality relation and gives another connection between the duality operation $\chi \rightarrow \chi^*$ and Jordan decomposition.

(2.8) THEOREM. *Let $s \in G$ be semisimple and $x \in G$ arbitrary. The quantity $\sum_{x \in \text{irr } G} \chi(x) \overline{\chi^*(s)}$ is equal to $|Z(s)|/\text{St}(s)$ if x_{ss} is conjugate to s and vanishes otherwise.*

Proof. Applying Corollary 2.5 we obtain

$$\sum_{x \in \text{irr } G} \chi(x) \overline{\chi^*(s)} = \text{St}(s)^{-1} \sum_{u \in V(s)} \sum_{x \in \text{irr } G} \chi(x) \overline{\chi(su)}.$$

An orthogonality formula for characters ([10, 31.19]) shows the quantity in question vanishes unless x is conjugate to some su , $u \in V(s)$, which occurs only if x_{ss} is conjugate to s . We may assume, therefore, that $x_{ss} = s$. From the above, $\sum_{x \in \text{irr } G} \chi(x) \overline{\chi^*(s)} = N |Z(x)|/\text{St}(s)$, where N is the number of su , $u \in V(s)$, conjugate to x . Since $N = |Z(s) : Z(x)|$, the theorem is proved.

(2.9) COROLLARY. *If $u \in G$ is unipotent, then $\sum_{x \in \text{irr } G} \chi(x) \chi^*(1) = |G|_{p'}.$ In particular, $\sum_{x \in \text{irr } G} \chi(1) \chi^*(1) = |G|_{p'}.$*

3. A SECOND METHOD

In this section we assume $G = \mathbf{G}^F$ is the group of rational points of a connected, reductive algebraic group \mathbf{G} defined over a finite field \mathbb{F}_q with Frobenius endomorphism F . The group $G = \mathbf{G}^F$ has a split (B, N) -pair of characteristic p , where $q = p^e$, with $B = \mathbf{B}_0^F$ for an F -stable Borel subgroup \mathbf{B}_0 of \mathbf{G} , $T = \mathbf{T}_0^F$ for an F -stable maximal torus $\mathbf{T}_0 \leq \mathbf{B}_0$, $N = N(\mathbf{T}_0)^F$ and $W = \mathbf{W}^F$ for $\mathbf{W} = N(\mathbf{T}_0)/\mathbf{T}_0$. Let $\text{rank}_q(\mathbf{G})$ be the dimension of a maximal \mathbb{F}_q -split torus in \mathbf{G} , and define $\varepsilon_{\mathbf{G}} = (-1)^r$, $r = \text{rank}_q(\mathbf{G})$.

If \mathbf{T} is an F -stable torus in \mathbf{G} , then $q^{-1}F$ acts on the character group $X(\mathbf{T})$ as an automorphism of finite order. Let $\lambda_{\mathbf{T}}(u)$ be the characteristic polynomial of $q^{-1}F$ on $X(\mathbf{T})$ ($\lambda_{\mathbf{T}}(u)$ has integer coefficients). The multiplicity of $u - 1$ as a factor of $\lambda_{\mathbf{T}}(u)$ is equal to $\text{rank}_q(\mathbf{T})$. Thus $u^d \lambda_{\mathbf{T}}(1/u) = \varepsilon_{\mathbf{T}} \lambda_{\mathbf{T}}(u)$, where $d = \dim \mathbf{T}$. Also, $\lambda_{\mathbf{T}}(q) = |\mathbf{T}^F|$. If $\mathbf{T}_0 \leq \mathbf{B}_0 \leq \mathbf{G}$ are as above, then

$|\mathbf{G}^F| = q^{|\Phi^+|} \lambda_{T_0}(q) P_G(q)$, where Φ is the root system of \mathbf{G} and $P_G(u) = \sum_{w \in W^F} u^{l(w)}$ is the Poincaré polynomial of (\mathbf{G}, F) .

Let \mathbf{T} be an F -stable maximal torus of \mathbf{G} , and let $\theta: \mathbf{T}^F \rightarrow C^*$ be a linear character. Deligne and Lusztig [11] have constructed, for each such pair (\mathbf{T}, θ) , a virtual character $R_{\mathbf{T}}^{\mathbf{G}} \theta$ of \mathbf{G}^F . Let $s \in \mathbf{G}^F$ be semisimple, and assume $\chi \in \text{char } \mathbf{G}^F$. Theorem 6.8 and Propositions 7.3 and 7.5 of [11] imply that

$$\chi(s) = \sum_{(\mathbf{T}, \theta)} \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{G}} \frac{(\chi, R_{\mathbf{T}}^{\mathbf{G}} \theta)}{(R_{\mathbf{T}}^{\mathbf{G}} \theta, R_{\mathbf{T}}^{\mathbf{G}} \theta)} \frac{\theta^{\mathbf{G}^F}(s)}{\text{St}(s)}, \quad (3.1)$$

where the sum is over the \mathbf{G}^F -conjugacy classes of pairs (\mathbf{T}, θ) .

If \mathbf{T} contains a \mathbf{G}^F -conjugate of s , define $f_{\mathbf{T}, s}(q) = |\mathbf{Z}(s)^{0F}|_p / |\mathbf{T}^F|$, interpreted as a polynomial (with integer coefficients) evaluated at q . Also, set

$$g_{\mathbf{T}, \theta, x, s} = \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{Z}(s)^0} |\mathbf{Z}(s)^F : \mathbf{Z}(s)^{0F}| \frac{(\chi, R_{\mathbf{T}}^{\mathbf{G}} \theta)}{(R_{\mathbf{T}}^{\mathbf{G}} \theta, R_{\mathbf{T}}^{\mathbf{G}} \theta)} \sum \theta t,$$

where $\sum \theta t$ is summed over all \mathbf{G}^F -conjugates t of s in \mathbf{T}^F . Since $\text{St}(s) = \varepsilon_{\mathbf{G}} \varepsilon_{\mathbf{Z}(s)^0} |\mathbf{Z}(s)^{0F}|_p$, (3.1) implies the following.

$$\chi(s) = \sum_{(\mathbf{T}, \theta)} g_{\mathbf{T}, \theta, x, s} f_{\mathbf{T}, s}(q). \quad (3.2)$$

The right-hand side of (3.2) can be interpreted as a polynomial (with complex coefficients) evaluated at q . Let \mathbf{H} be an F -stable maximal torus contained in an F -stable Borel subgroup of $\mathbf{Z}(s)^0$. Then

$$\begin{aligned} \text{St}(s) f_{\mathbf{T}, s}(1/q) &= \text{St}(s) \lambda_{\mathbf{H}}(1/q) P_{\mathbf{Z}(s)^0}(1/q) \lambda_{\mathbf{T}}(1/q)^{-1} \\ &= \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{G}} \lambda_{\mathbf{H}}(q) P_{\mathbf{Z}(s)^0}(q) \lambda_{\mathbf{T}}(q)^{-1} \\ &= \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{G}} f_{\mathbf{T}, s}(q), \end{aligned}$$

since $\dim \mathbf{H} = \dim \mathbf{T}$. We now need the following result, first communicated to the author by G. Lusztig (private communication).

(3.3) THEOREM (Lusztig). *If \mathbf{T} is an F -stable maximal torus and $\theta: \mathbf{T}^F \rightarrow C^*$ is a linear character, then $(R_{\mathbf{T}}^{\mathbf{G}} \theta)^* = \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{G}} R_{\mathbf{T}}^{\mathbf{G}} \theta$.*

Combining Theorem 3.3 with the observations above, we have proved the next theorem.

(3.4) THEOREM. *Let $s \in \mathbf{G}^F$ be semisimple and let $\chi \in \text{char } \mathbf{G}^F$. Then $\chi^*(s) = \text{St}(s) \chi^{\#}(s)$, where $\chi^{\#}(s)$ is obtained from expression (3.2) for $\chi(s)$ by replacing q by $1/q$.*

Now consider the case $s = 1$. Formula (3.2) takes the following form (see Lusztig [19, 3.14]).

$$\chi(1) = \sum_{(\mathbf{T}, \theta)} \varepsilon_{\mathbf{T}} \varepsilon_{\mathbf{G}} \frac{(\chi, R_{\mathbf{T}}^{\mathbf{G}} \theta)}{(R_{\mathbf{T}}^{\mathbf{G}} \theta, T_{\mathbf{T}}^{\mathbf{G}} \theta)} \frac{|\mathbf{G}^F|_{p'}}{|\mathbf{T}^F|}. \quad (3.5)$$

The sum in (3.5) is over the \mathbf{G}^F -classes of pairs (\mathbf{T}, θ) . The right-hand side of (3.5) is viewed as the value $d_{\chi}(q)$ of a polynomial $d_{\chi}(u)$ with rational coefficients ($\chi \in \text{char } \mathbf{G}^F$).

(3.6) COROLLARY. *If $\chi \in \text{char } \mathbf{G}^F$, then $d_{\chi}(q) = q^{|\Phi^+|} d_{\chi}(1/q)$, where Φ is the root system of \mathbf{G} .*

Let Φ be a fixed root system. If we consider all connected, reductive \mathbf{G} defined over finite fields with root system Φ , then the resulting set of polynomials $d_{\chi}(u)$, $\chi \in \text{irr } \mathbf{G}^F$, is finite [11, Proposition 6.3 and Theorem 6.8]. If we further assume \mathbf{G} is semisimple, then there are integers M , N , and E (depending only on Φ) such that $|\mathbf{G}^F|$ always divides $q^M(q^N - 1)^E$ (see Steinberg [23, 11.16]). Also, if $\chi \in \text{irr } \mathbf{G}^F$, then $\chi(1)$ divides $|\mathbf{G}^F|$ [10, 33.7]. Thus if we assume q is large (with respect to Φ) and \mathbf{G} is semisimple, the polynomials $d_{\chi}(u)$, $\chi \in \text{irr } \mathbf{G}^F$, can be factored as $d_{\chi}(u) = \alpha u^e \phi(u)$ with α rational, e a nonnegative integer and $\phi(u)$ a product of cyclotomic polynomials. This fact, together with Corollary 3.6, proves the following.

(3.7) COROLLARY. *Let Φ be a root system. Assume \mathbf{G} is a semisimple group defined over F_q , with root system Φ and Frobenius endomorphism F . If q is large with respect to Φ , and if $\chi \in \text{irr } \mathbf{G}^F$, then $\chi^*(1)/\chi(1)$ is of the form $\pm q^n$ for some integer n .*

4. DUALITY AND CHARACTER DEGREES

From now on, \mathbf{G} will denote an algebraic group which is defined over a finite field \mathbb{F}_q , with corresponding Frobenius endomorphism F . It will be assumed that \mathbf{G} is at least connected and reductive. Let $V(1)$ be the set of unipotent element of \mathbf{G}^F . We will consider the following two statements about the pair (\mathbf{G}, F) .

(4.1) For any $\chi \in \text{irr } \mathbf{G}^F$, the quantity $C(\chi) = \sum_{u \in V(1)} \chi(u)/\chi(1)$ is of the form $\pm q^e$ for some nonnegative integer e .

(4.2) For any $\chi \in \text{irr } \mathbf{G}^F$, the quantity $\chi^*(1)/\chi(1)$ is of the form $\pm q^n$ for some integer n .

For $\chi \in \text{irr } \mathbf{G}^F$, let $\chi' \in \text{irr } \mathbf{G}^F$ be such that $\chi^* = \pm \chi'$ (Theorem 1.5). Then $C(\chi)$ and $C(\chi')$ are algebraic integers [10, p. 236]. Also, $C(\chi) = |\mathbf{G}^F|_p \chi^*(1)/\chi(1)$ and $C(\chi') = |\mathbf{G}^F|_p \chi(1)/\chi^*(1)$ by Corollary 2.6, since $\chi^{**} = \chi$ (Theorem 1.5). Therefore $C(\chi)$ is a rational integer, and $\chi(1)_{p'} = |\chi^*(1)|_{p'}$. Since $|\mathbf{G}^F|_p = q^{|\Phi^+|}$, where Φ is the root system of \mathbf{G} , we have proved the following.

(4.3) PROPOSITION. *The pair (\mathbf{G}, F) satisfies (4.1) if and only if (\mathbf{G}, F) satisfies (4.2).*

By Corollary 3.7, we know that (4.1) and (4.2) are valid for (\mathbf{G}, F) if \mathbf{G} is semisimple and q is sufficiently large with respect to Φ .

The proof of the next lemma is straightforward and will be omitted.

(4.4) LEMMA. *Let $G = \langle B, N \rangle$ be a finite group with split (B, N) -pair of characteristic p , and assume H is a subgroup of $T \cap Z(G)$, $T = B \cap N$. Let $\bar{G} = G/H$, and give \bar{G} the natural induced split (B, N) -pair structure. For $\chi \in \text{char } \bar{G}$, let $L\chi \in \text{char } G$ be the lift of χ to G via the projection $G \rightarrow \bar{G}$, so that $L\chi(g) = \chi(gH)$. Then $(L\chi)^* = L(\chi^*)$. Thus if $\chi \in \text{irr } \bar{G}$, $\chi^*(1)/\chi(1) = (L\chi)^*(1)/L\chi(1)$.*

(4.5) LEMMA. *Suppose that G and G' are finite groups with split (B, N) -pairs of characteristic p , that $G \trianglelefteq G'$ and that G and G' have a common Coxeter system (W, R) . For $J \subseteq R$ let P_J (P'_J) be the corresponding standard parabolic subgroup of G (G' , respectively). Assume $O_p(P_J) = O_p(P'_J)$ for $J \subseteq R$. Let $\chi \in \text{irr } G'$, and let $\chi_0 \in \text{irr } G$ be any component of the restriction $\chi|_G$ of χ to G . Then $\chi^*(1)/\chi(1) = \chi_0^*(1)/\chi_0(1)$.*

Proof. First note that

$$\chi^*(1) = \sum_{J \subseteq R} (-1)^{|J|} |G' : P'_J| (\chi|_{V_J}, 1_{V_J})_{V_J},$$

where $V'_J = O_p(P'_J)$. Also, $V'_J = O_p(P_J) = V_J$ for $J \subseteq R$. Thus $U = V_\phi = V'_\phi = U'$, so that G contains all p -elements of G' . If $J \subseteq R$, then $GP'_J = P'_K$ for some $K \subseteq R$. Then $V'_K \subseteq O_p(G) = V_R = \{1\}$, and therefore $K = R$ and $GP'_J = G'$. Also, $P_J = N_G(V_J) = G \cap N_{G'}(V'_J)$ (Curtis [8, Proposition 1.5]), and therefore $|G : P_J| = |G' : P'_J|$, for $J \subseteq R$. It follows that $\chi^*(1) = (\chi|_G)^*(1)$.

By Clifford's theorem ([10, 49.7]) we have $\chi|_G = m(\chi_0 + \chi_1 + \cdots + \chi_{k-1})$, where $m \geq 1$ and $\chi_0, \dots, \chi_{k-1}$ are distinct G' -conjugate irreducible characters of G . Thus $\chi_0(1) = \cdots = \chi_{k-1}(1)$ and $\chi(1) = mk\chi_0(1)$. Since $G \trianglelefteq G'$, the set $V(1)$ of unipotent elements of G is G' -invariant. Therefore $\sum_{u \in V(1)} \chi_0(u) = \cdots = \sum_{u \in V(1)} \chi_{k-1}(u)$. From this it follows that $\chi_0^*(1) = \cdots =$

$\chi_{k-1}^*(1)$ by Corollary 2.6. Therefore $\chi^*(1) = (\chi|_G)^*(1) = mk\chi_0^*(1)$, and the proof of the lemma is completed.

Let $\pi: \tilde{\mathbf{G}} \rightarrow \mathbf{G}$ be an isogeny of semisimple groups. Assume $\tilde{\mathbf{G}}, \mathbf{G}$ are defined over \mathbb{F}_q , with Frobenius endomorphisms \tilde{F}, F , and that $\pi\tilde{F} = F\pi$ (i.e., π is defined over \mathbb{F}_q). Let $\tilde{G} = \tilde{\mathbf{G}}^{\tilde{F}}$, $G = \mathbf{G}^F$ and $\tilde{H} = (\ker \pi)^{\tilde{F}}$. The next lemma follows from Lemmas 4.4 (applied to $\tilde{G} \rightarrow \tilde{G}/\tilde{H}$) and 4.5 (applied to $\tilde{G}/\tilde{H} \cong \pi(\tilde{G}) \trianglelefteq G$).

(4.6) LEMMA. *Assume $\pi: \tilde{\mathbf{G}} \rightarrow \mathbf{G}$ is an isogeny of semisimple groups defined over \mathbb{F}_q . If (4.2) holds for $(\tilde{\mathbf{G}}, \tilde{F})$, then (4.2) holds for (\mathbf{G}, F) .*

(4.7) LEMMA. *Assume that \mathbf{G} is connected, reductive and defined over \mathbb{F}_q , with Frobenius endomorphism F . Let \mathbf{G}_0 be a closed, connected, reductive F -stable subgroup of \mathbf{G} , and assume $D\mathbf{G}_0 = D\mathbf{G}$. Then (4.2) holds for (\mathbf{G}, F) if and only if (4.2) holds for (\mathbf{G}_0, F) .*

Proof. Apply Lemma 4.5 to $\mathbf{G}_0^F \trianglelefteq \mathbf{G}^F$.

We now show that (4.2) will hold in general if (4.2) is true for certain special cases (see Lusztig [16, 1.18] for a similar reduction argument). To prove that (4.2) is true for (\mathbf{G}, F) , it is enough to prove (4.2) holds for $(D\mathbf{G}, F)$ (Lemma 4.7). Thus we may assume $\mathbf{G} = D\mathbf{G}$ is semisimple. Let $\mathbf{G}_1, \dots, \mathbf{G}_m$ be the simple components of \mathbf{G} (see Humphreys [14, 27.5]). Let $\mathbf{H}_1, \dots, \mathbf{H}_k$ be the subgroups of \mathbf{G} corresponding to the orbits of F on $\{\mathbf{G}_1, \dots, \mathbf{G}_m\}$. Then $F\mathbf{H}_i = \mathbf{H}_i$, and multiplication $\mathbf{H}_1 \times \dots \times \mathbf{H}_k \rightarrow \mathbf{G}$ is an isogeny defined over \mathbb{F}_q . By Lemma 4.6, we may reduce to the case $\mathbf{G} = \mathbf{H}_1$, and may assume $F\mathbf{G}_i = \mathbf{G}_{i+1}$ ($1 \leq i < m$), $F\mathbf{G}_m = \mathbf{G}_1$. Let $\tilde{\mathbf{G}} = \mathbf{G}_1 \times \dots \times \mathbf{G}_m$, and define $\tilde{F}: \tilde{\mathbf{G}} \rightarrow \tilde{\mathbf{G}}$ by $\tilde{F}(g_1, \dots, g_m) = (Fg_m, Fg_1, \dots, Fg_{m-1})$. Then \tilde{F} is the Frobenius endomorphism for an \mathbb{F}_q -rational structure on $\tilde{\mathbf{G}}$, and multiplication $\tilde{\mathbf{G}} \rightarrow \mathbf{G}$ is an isogeny defined over \mathbb{F}_q . Applying Lemma 4.6 again, we may assume $\mathbf{G} = \mathbf{G}_1 \times \dots \times \mathbf{G}_m$ and that $F\mathbf{G}_i = \mathbf{G}_{i+1}$ ($1 \leq i < m$), $F\mathbf{G}_m = \mathbf{G}_1$. Note that \mathbf{G}_1 is defined over \mathbb{F}_{q^m} , with Frobenius endomorphism F^m . Also, $\mathbf{G}_1^{F^m} \cong \mathbf{G}^F$. Therefore (4.2) holds for (\mathbf{G}, F) if (4.2) holds for (\mathbf{G}_1, F^m) (with q^m replacing q). We thus reduce further to the case $\mathbf{G} = \mathbf{G}_1$, so that the Dynkin diagram Γ for \mathbf{G} is connected. Applying Lemma 4.6 once more, we may also assume that \mathbf{G} is simply connected. The reduction is summarized in the following theorem.

(4.7) THEOREM. *Let $\Gamma_1, \dots, \Gamma_k$ be connected Dynkin diagrams, and let q be a prime power. Assume that (4.2) holds for $(\tilde{\mathbf{G}}, \tilde{F})$ whenever $\tilde{\mathbf{G}}$ is a simply connected group defined over \mathbb{F}_{q^m} , $m \geq 1$, with Frobenius endomorphism \tilde{F} and Dynkin diagram Γ_i for some i . Let \mathbf{G} be a connected, reductive group defined over \mathbb{F}_q with Frobenius endomorphism F , such that the components of the Dynkin diagram Γ of \mathbf{G} are among $\Gamma_1, \dots, \Gamma_k$. Then (4.2) holds for (\mathbf{G}, F) .*

(4.8) THEOREM. Let G be simply connected and defined over \mathbb{F}_q , with Frobenius endomorphism F . Assume (G, F) is of type $A_n, {}^2A_n, B_n, C_n, D_n$ or 2D_n . Then (4.2) holds for (G, F) .

Proof. By Lemma 4.7, it is sufficient to show (4.2) holds when G is reductive, DG is simply connected, $Z(G)$ is connected, and (G, F) is of type $A_n, {}^2A_n, B_n, C_n, D_n$ or 2D_n . Let $W = W^F$ be the (relative) Weyl group of $G = G^F$, with set of distinguished generators R . Suppose $\chi \in \text{irr } G$, and let $\chi' \in \text{irr } G$ be such that $\chi' = \pm \chi^*$ (Theorem 1.5). By the remarks after Theorem 1.5, χ and χ' are both components of some induced cuspidal representation $I_J \varphi$, $J \subseteq R$. Let $W(\varphi)$ be the set of distinguished (W_J, W_J) -double coset representatives w such that ${}^w J = J$ and ${}^w \varphi = \varphi$. Thus $W(\varphi)$ is isomorphic to the stabilizer of φ in $N_w(W_J)/W_J$. Also, $W(\varphi)$ is a Coxeter group, with set of distinguished generators S , and the irreducible components of $I_J \varphi$ correspond by specialization to the irreducible characters of $W(\varphi)$ (see Asai [4, Sect. 7]). Let χ correspond to μ and χ' to μ' , $\mu, \mu' \in \text{irr } W(\varphi)$. Assume $|S| \geq 1$ (if $W(\varphi) = \{1\}$, then $\chi = \chi' = \pm \chi^*$; so $\chi^*(1)/\chi(1) = \pm 1$ and the assertion of (4.2) holds for (G, F, χ)). We shall prove $\mu' = \text{sgn} \otimes \mu$, where sgn is the sign character of $(W(\varphi), S)$.

Assume $|S| = 1$, so that $|W(\varphi)| = 2$ and $I_J \varphi = \chi + \chi_1$, $\chi_1 \in \text{irr } G$, $\chi_1 \neq \chi$. We must show $\chi' = \chi_1$. By Curtis's result (Theorem 1.4), Lemma 1.2 and [4, 1.3.18], we may assume J is a maximal subset of R . Then by [8, Theorem 3.5], we have $\chi^* = (-1)^{|R|-1} m I_J \varphi + (-1)^{|R|} \chi$, for some $m > 0$. Thus $m(\chi + \chi_1) = \pm \chi \pm \chi^*$, and so $\chi_1 = \chi' = \pm \chi^*$ since $\chi_1 \neq \chi$. It follows that $\mu' \neq \mu$, and therefore $\mu' = \text{sgn} \otimes \mu$ in this case.

Assume that $|S| > 1$. By a suitable induction hypothesis, Theorem 1.4 and Asai [4, Theorem 7.2.5], we may assume that $(\mu', \text{Ind}_{W(\varphi)_K}^{W(\varphi)} \nu) = (\text{sgn} \otimes \mu, \text{Ind}_{W(\varphi)_K}^{W(\varphi)} \nu)$ when $K \subseteq S$, $K \neq S$, and $\nu \in \text{irr } W(\varphi)_K$, where $W(\varphi)_K$ is generated by K . It follows that $\mu' = \text{sgn} \otimes \mu$ (see Benson and Curtis [5]), since $|S| > 1$.

The irreducible characters of $G = G^F$ are partitioned into geometric conjugacy classes $\text{irr}(G^F, [s])$, one for each conjugacy class $[s]$ of semisimple elements of G^{*F} , where G^* is the dual group of G (Deligne and Lusztig [11, Sects. 5 and 6]). Let $\chi \in \text{irr}(G^F, [s])$. Then $\chi' \in \text{irr}(G^F, [s])$ by the remarks above and Asai [4, Theorem 6.1]. Let $R: \text{irr}(G^F, [s]) \rightarrow \text{irr}(Z_{G^*}(s)^{*F}, [1])$ be the bijection defined in Lusztig [18] and Asai [4], and set $\rho = R(\chi)$, $\rho' = R(\chi')$. Then $\chi'(1)/\chi(1) = \rho'(1)/\rho(1)$ (see [18, Theorem 8.2; 4, Theorem 6.1]). Also, ρ and ρ' are both components of the character induced from some irreducible cuspidal unipotent character ρ_1 of some Levi factor of $Z_{G^*}(s)^{*F}$. By Lusztig [19, 3.26.1], $\rho'(1)/\rho(1) = d(\chi_{\mu'})/d(\chi_{\mu})$, where $d(\chi_{\mu})$ and $d(\chi_{\mu'})$ are specializations of the generic degrees for the characters of the generalized Hecke algebra of $I_J \varphi$ which correspond to μ and μ' . Since $\mu' = \text{sgn} \otimes \mu$, $d(\chi_{\mu'})$ can be obtained from $d(\chi_{\mu})$ by replacing q by $1/q$ and multiplying by a power of q (see Green [12]). Hence $d(\chi_{\mu'})/d(\chi_{\mu})$ is of the

form q^n for some integer n , by the formulas of Hoefsmit [13]. Therefore $\chi^*(1)/\chi(1) = \pm\chi'(1)/\chi(1) = \pm\rho'(1)/\rho(1) = \pm q^n$; so that (4.2) holds for G, F, χ . This completes the proof of the theorem.

If the characteristic p of \mathbb{F}_q is not 2, then a simpler proof of Theorem 4.8 is possible. Let G be as in the proof above. The formulas of Lusztig [18, 19], Asai [4], and Hoefsmit [13] show that $\chi(1)_p$ is always a power of q when $\chi \in \text{irr } G^F$. Thus $\chi^*(1)/\chi(1)$ is trivially of the form $\pm q^n$ in this case, since $\pm\chi^* \in \text{irr } G^F$.

If G is simply connected and (G, F) is of type ${}^3D_4, E_6, {}^2E_6, E_7, E_8, F_4$ or G_2 , then (4.2) holds for (G, F) provided q is sufficiently large (Corollary 3.7). Combining this with Theorems 4.7 and 4.8 proves the following.

(4.9) THEOREM. *Assertion (4.2) holds for any (G, F) , provided q is large.*

Note that the sense in which q must be large in Theorem 4.9 does not depend on the particular root system of G (as in Corollary 3.7). It is expected that (4.2) is true in general.

ACKNOWLEDGMENT

The author wishes to thank C. W. Curtis for much help with [3] and George Lusztig for several helpful suggestions.

REFERENCES

1. D. ALVIS, The duality operation in the character ring of a finite Chevalley group, *Bull. Amer. Math. Soc.* **1** (1979), 907–911.
2. D. ALVIS, Duality in the character ring of a finite Chevalley group, in "Proc. Symp. pure Math." No. 37, pp. 353–357, Amer. Math. Soc., Providence, R.I., 1980.
3. D. ALVIS, "Duality and Its Applications to the Character Theory of Finite Lie Groups," Ph.D. dissertation, University of Oregon, Eugene, 1980.
4. T. ASAI, Endomorphism algebras of the reductive groups over F_q of classical type, preprint.
5. C. T. BENSON AND C. W. CURTIS, On the degrees and rationality of certain characters of finite Chevalley groups, *Trans. Amer. Math. Soc.* **165** (1972), 251–273; **202** (1975), 405–406.
6. R. CARTER AND G. LUSZTIG, Modular representations of finite groups of Lie type, *Proc. London Math. Soc.* (3) **32** (1976), 347–384.
7. C. W. CURTIS, The Steinberg character of a finite group with a (B, N) -pair, *J. Algebra* **4** (1966), 433–441.
8. C. W. CURTIS, Reduction theorems for characters of finite groups of Lie type, *J. Math. Soc. Japan* **27** (1975), 666–688.

9. C. W. CURTIS, Truncation and duality in the character ring of a finite group of Lie type, *J. Algebra* **62** (1980), 320–332.
10. C. W. CURTIS AND I. REINER, "Representation Theory of Finite Groups and Associative Algebras," Interscience, New York, 1962.
11. P. DELIGNE AND G. LUSZTIG, Representations of reductive groups over finite fields, *Ann. of Math.* (2) **103** (1976), 103–161.
12. J. A. GREEN, On the Steinberg characters of finite Chevalley groups, *Math. Z.* **117** (1970), 272–288.
13. P. N. HOEFSMIT, "Representations of Hecke Algebras of Finite Groups with BN -pairs of Classical Type," Ph.D. dissertation, University of British Columbia, Vancouver, 1974.
14. J. E. HUMPHREYS, "Linear Algebraic groups," Springer-Verlag, New York, 1975.
15. N. KAWANAKA, Unipotent elements and characters of finite Chevalley groups, *Osaka J. Math.* **12** (1975), 523–554.
16. G. LUSZTIG, Coxeter orbits and eigenspaces of Frobenius, *Invent. Math.* **38** (1976), 101–159.
17. G. LUSZTIG, Divisibility of projective modules of finite Chevalley groups by the Steinberg module, *Bull. London Math. Soc.* **8** (1976), 130–134.
18. G. LUSZTIG, Irreducible representations of finite classical groups, *Invent. Math.* **43** (1977), 125–175.
19. G. LUSZTIG, Representations of finite Chevalley groups, in "C.B.M.S. Regional Conf. Ser. in Math. No. 39," Amer. Math. Soc., Providence, R.I., 1978.
20. F. A. RICHEN, Modular representations of split (B, N) -pairs, *Trans. Amer. Math. Soc.* **140** (1969), 435–460.
21. T. A. SPRINGER, Cusp forms for finite groups, Part C, in "Seminar on Algebraic Groups and Related Finite Groups," Lecture Notes in Math. No. 131, Springer-Verlag, New York, 1970.
22. T. A. SPRINGER, A formula for the characteristic function on the unipotent set of a finite Chevalley group, *J. Algebra* **62** (1980), 393–399.
23. R. STEINBERG, Endomorphisms of linear algebraic groups, *Mem. Amer. Math. Soc.*, No. 80 (1968).